## ARTHROSCOPIC BANKART REPAIR

## This protocol to be used for Laterjet procedures as well

## Dr. Troy Berg, Dr. Nathan Harris

Patient\_\_\_\_\_ D.O.S.\_\_\_\_\_

Chippewa Valley Orthopedics & Sports Medicine

1200 OakLeaf Way, Suite A Altoona, WI 54720

757 Lakeland Drive, Suite B Chippewa Falls, WI 54729

	WEEK 1-3 BEGINS DOS	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9-12	MONTH 4
PASSIVE ER At modified neutral	0-10° supine	20°	30°	30° Increase 10° /week until within 10° of opposite side, no strer weeks post-operatively				
ACTIVE/ AAROM ER	none	0	0	10°	20°	30°	40°	Increase 10°/wk until FULL
PASSIVE FLEXION	Supine 0-70° as tolerated	90° Increase as tolerated						FULL
ACTIVE/AA FLEXION	none	60° Increase as tolerated						FULL
IMMOB/ SLING	yes	yes	Wean as tolerated					
PT visits/wk	0	2	2	2	2	2	2	→ 2 1
EXERCISES	AROM (cervical elbow,wrist,hand) PROM of shoulder progressing to AAROM   PENDULUMS-gentle dangle, then intiate ROM and increase movement after week 2							ADVANCED PRE PNF, overhead ADVANCED SCAPULAR
Any questions,	JOINT MOBS (grade I & II )							
please contact :		AAROM FLEX/ER (swiss ball, wand, seated, pulleys) Progress to active						- STABIL.IZATION - Pushup+, Press-up,
Northwoods Therapy Associates	SCAP MOBS/STAB (swiss ball, reach & roll to chest, ret/dep, serratus, prone, CKC)							reverse corner press Side lying shoulder,
ALtoona, WI		ISOMETRICS (ER, FLEX,EXT,ABD)						
(715) 839–9266	UBE							
Chippewa Falls, WI (715) 723–6060 Reviewed	Follow specific MD instruction. Individual progression may vary.implicationAt 4-6 months initiate return to advance level tasks and repetitiveTur					PRE's below shoulder height, with improved scapular control move to overhead movement Tubing, dumbbells		PLYOMETRICS As appropriate for functional tasks
June 2018	overhead activity.						ATED ROW	