Proximal Humerus Fracture/Greater Tuberosity ORIF

Dr. Austin Crow, Dr. Evan Peissig

Chippewa Valley Orthopedics & Sports Medicine 1200 OakLeaf Way, Suite A 757 Lakeland Drive, Suite B		Patient DOS	
Altoona, WI 54720	Chippewa Falls, WI 54729		
	0-6 Weeks	6 Weeks – 3 Months	3-6 Months +
PT Visits	2x/wk	2-3x/wk	1-2x/wk, discharge as appropriate
HEP	Everyday	5x/wk	4-5x/wk
Sling	Wear at night 0-6 weeks	Discontinue	
	Wean during day at 2 weeks		
Exercises	PROM & AROM of neck, elbow, wrist, hand (5x/day)	Continue PROM in all planes Supine Stick ER, Scaption	Resisted Rows, Extension, IR, ER (tubing, dumbbells)
	Gentle passive pendulum (3x/day) Gentle PROM in all planes	Table Flexion, ER, Abd Pulleys	"W" Tubing Resisted PNF D1 & D2 (tubing, dumbbells)
		Swiss Ball Roll	Wall Push Ups, Push up Plus
	ICE: 3-5x/day for 30 minutes after exercises PRECAUTION: Avoid any active shoulder motion weeks 0-4	Standing Stick Flexion, ER, Abd	Advance strengthening as tolerated, progressing to normal work/sport
		Prone Extension, Rows Scaption, Abd	activities when appropriate
		Isometrics (Flexion, Extension, Abd, Add, IR, ER)	
		SA Punches	
		— Sideling ER	
		Scapular Retraction	
		Rhythmic Stabs	
		Shoulder Shrugs	
		Standing "I", "Y", "T"	