

# ARTHROSCOPIC BANKART REPAIR

This protocol to be used for Laterjet procedures as well

Dr. Troy Berg

Chippewa Valley Orthopedics & Sports Medicine

1200 OakLeaf Way, Suite A

Altoona, WI 54720

757 Lakeland Drive, Suite B

Chippewa Falls, WI 54729

Patient\_\_\_\_\_

D.O.S.\_\_\_\_\_

	WEEK 1-3 BEGINS DOS	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9-12	MONTH 4
<b>PASSIVE ER</b> At modified neutral	0-10° supine	20°	30°	Increase 10° /week until within 10° of opposite side, no strenuous ER until 10 weeks post-operatively				
<b>ACTIVE/ AAROM ER</b>	none	0	0	10°	20°	30°	40°	Increase 10°/wk until FULL
<b>PASSIVE FLEXION</b>	Supine 0-70° as tolerated	90°	Increase as tolerated					FULL
<b>ACTIVE/AA FLEXION</b>	none	60°	Increase as tolerated					FULL
<b>IMMOB/ SLING</b>	yes	yes	Wean as tolerated					
<b>PT visits/wk</b>	0	2	2	2	2	2	2	→ 2 1
<b>EXERCISES</b>	AROM (cervical elbow,wrist,hand)	PROM of shoulder progressing to AAROM						<b>ADVANCED PRE</b> PNF, overhead
	PENDULUMS-gentle dangle, then initiate ROM and increase movement after week 2							<b>ADVANCED SCAPULAR STABILIZATION</b> Pushup+, Press-up, reverse corner press Side lying shoulder, reach & roll-full
<b>Any questions, please contact : Northwoods Therapy Associates Altoona, WI (715) 839-9266 Chippewa Falls, WI (715) 723-6060</b>	JOINT MOBS (grade I & II )		AAROM FLEX/ER (swiss ball, wand, seated, pulleys) Progress to active					
	SCAP MOBS/STAB (swiss ball, reach & roll to chest, ret/dep, serratus, prone, CKC)							
	ISOMETRICS (ER, FLEX,EXT,ABD)							
	UBE							
Reviewed June 2018	Follow specific MD instruction. Individual progression may vary. At 4-6 months initiate return to advance level tasks and repetitive overhead activity.					PRE's below shoulder height, with improved scapular control move to overhead movement Tubing, dumbbells		<b>PLYOMETRICS</b> As appropriate for functional tasks
SEATED ROW								